


|  |                                    |                                 |
|--|------------------------------------|---------------------------------|
|  | <h1>NCAM1 Polyclonal Antibody</h1> | E<br>9<br>1<br>2<br>1<br>1<br>2 |
|--|------------------------------------|---------------------------------|

|                           |   |
|---------------------------|---|
| <b>Swiss-Prot No.:</b>    | P13591  |
| <b>Storage/Stability:</b> | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.   |
| <b>Immunogen:</b>         | Recombinant protein of human NCAM1 (NP_851996.2).   |
| <b>Purification:</b>      | Affinity purification   |
| <b>Reactivity:</b>        | Human,Mouse   |
| <b>Other Names:</b>       | —   |
| <b>Background:</b>        | This gene encodes a cell adhesion protein which is a member of the immunoglobulin superfamily. The encoded protein is involved in cell-to-cell interactions as well as cell-matrix interactions during development and differentiation. The encoded protein has been shown to be involved in development of the nervous system, and for cells involved in the expansion of T cells and dendritic cells which play an important role in immune surveillance. Alternative splicing results in multiple transcript variants. |
| <b>Gene ID:</b>           | 4684  |
| <b>Source:</b>            | Rabbit  |
| <b>Antibody type:</b>     | Polyclonal Antibodies   |
| <b>Isotype:</b>           | IgG   |
| <b>Molecular</b>          | 40kDa/73kDa/80kDa/83kDa/93kDa/94kDa   |

**For Research Use Only**

Order: [order@enogene.com](mailto:order@enogene.com) Support: [service@enogene.com](mailto:service@enogene.com) Web:  
[www.enogene.com](http://www.enogene.com)

---

|                               |                   |
|-------------------------------|-------------------|
| <b>Weight:</b>                |                   |
| <b>Recommended Dilutions:</b> | WB 1:500 - 1:2000 |

---

**For Research Use Only**