

	VEGF Polyclonal Antibody	E 9 1 2 3 0 3
--	-----------------------------------	---------------------------------

Swiss-Prot No.:	P15692
Storage/Stability:	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Immunogen:	Recombinant protein of human VEGFA
Purification:	Affinity purification
Reactivity:	Human, Mouse, Rat
Other Names:	—
Background:	This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome, also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended

For Research Use Only

	isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site.
Gene ID:	7422
Source:	Rabbit
Antibody type:	Polyclonal Antibodies
Isotype:	IgG
Molecular Weight:	15-27kDa/34-45kDa
Recommended Dilutions:	WB 1:500 - 1:2000 IF 1:50 - 1:200