

| | | |
|--|---|---|
|  | <h1 style="text-align: center;">SND1</h1> | <div style="background-color: #cccccc; padding: 5px; text-align: center;">E 9 5 8 7 4</div> |
|--|---|---|

| | |
|---------------------------|---|
| Antibody type: | Polyclonal Antibody |
| Applications: | WB IHC IF |
| Reactivity: | Human Mouse Rat |
| Molecular Weight: | 102kDa |
| Immunogen: | Recombinant protein of human SND1 |
| Gene ID: | 27044 |
| Swiss-Prot No.: | Q7KZF4 |
| Altername: | TDRD11;Tudor-SN;p100 |
| Source: | Rabbit |
| Isotype: | IgG |
| Purification: | Affinity purification |
| Storage/Stability: | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3. |
| Background: | This gene encodes a transcriptional co-activator that interacts with the acidic domain of Epstein-Barr virus nuclear antigen 2 (EBNA 2), a transcriptional activator that is required for B-lymphocyte transformation. Other transcription factors that interact with this protein are signal transducers and activators of transcription, STATs. This protein is also thought to be essential for normal cell growth. A similar protein in mammals and other organisms is a component of |

For Research Use Only

| | |
|-------------------------------|--|
| | the RNA-induced silencing complex (RISC). |
| Dilution: | WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200 |
| Shipping&Stablity: | Aliquot and store at -20°C. Avoid repeated freeze / thaw cycles. |