



<b>Antibody type:</b>	Polyclonal Antibody
<b>Applications:</b>	WB
<b>Reactivity:</b>	Human
<b>Molecular Weight:</b>	77kDa
<b>Immunogen:</b>	A phospho specific peptide corresponding to residues surrounding T638/641 of human PKCalpha/beta II
<b>Gene ID:</b>	5578
<b>Swiss-Prot No.:</b>	P17252
<b>Altename:</b>	AAG6;PKC-alpha;PKCA;PRKACA
<b>Source:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Purification:</b>	Affinity purification
<b>Storage/Stability:</b>	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
	Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol

<b>Background:</b>	esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This kinase has been reported to play roles in many different cellular processes, such as cell adhesion, cell transformation, cell cycle checkpoint, and cell volume control. Knockout studies in mice suggest that this kinase may be a fundamental regulator of cardiac contractility and Ca(2+) handling in myocytes.
<b>Dilution:</b>	WB 1:500 - 1:2000
<b>Shipping&amp;Stability:</b>	Aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.