

Immunotag™ HDAC7 Antibody

Antibody Specification	
Catalog No.	ITA0118
Product Description	Immunotag™ HDAC7 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	HDAC7
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC,IF/ICC,ELISA
Recommended Dilution	WB: 1:500~1:3000 IHC: 1:50~1:200, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human HDAC7
Specificity	HDAC7 antibody detects endogenous levels of total HDAC7
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	HDAC7
Accession No.	Q8WUI4
Alternate Names	DKFZP586J0917; FLJ99588; HD 7a; HD7; HD7a; HDAC 7; HDAC 7A; Hdac7; HDAC7_HUMAN; HDAC7A; Histone deacetylase 7; Histone deacetylase 7A; OTTHUMP00000202813; OTTHUMP00000202814;

Antibody Specification

Description	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Involved in muscle maturation by repressing transcription of myocyte enhancer factors such as MEF2A, MEF2B and MEF2C. During muscle differentiation, it shuttles into the cytoplasm, allowing the expression of myocyte enhancer factors (By similarity). May be involved in Epstein-Barr virus (EBV) latency, possibly by repressing the viral BZLF1 gene. Positively regulates the transcriptional repressor activity of FOXP3 (PubMed:17360565).
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	105kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.