

Immunotag™ MED17 Antibody

Antibody Specification	
Catalog No.	ITA0342
Product Description	Immunotag™ MED17 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	MED17
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF/ICC,ELISA
Recommended Dilution	WB 1:500-1:2000, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human MED17
Specificity	MED17 Antibody detects endogenous levels of MED17
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	MED17
Accession No.	Q9NVC6

Antibody Specification

Alternate Names	77-KD subunit; Activator recruited cofactor 77 kDa component; Activator-recruited cofactor 77 kDa component; ARC77; Cofactor required for Sp1 transcriptional activation subunit 6; cofactor required for Sp1 transcriptional activation, subunit 6, 77kDa; CRSP complex subunit 6; CRSP6; CRSP77; DRIP80; med17; MED17_HUMAN; Mediator complex subunit 17; Mediator of RNA polymerase II transcription subunit 17; Mediator of RNA polymerase II transcription subunit 17, S cerevisiae, homolog of; Thyroid hormone receptor-associated protein complex 80 kDa component; Thyroid hormone receptor-associated protein, 80-KD subunit; Transcriptional coactivator CRSP77; Trap80; VITAMIN D RECEPTOR-INTERACTING PROTEIN, 80-KD; Vitamin D3 receptor-interacting protein complex 80 kDa component;
Description	Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	73kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.