

Immunotag™ XRN2 Antibody

Antibody Specification	
Catalog No.	ITA0372
Product Description	Immunotag™ XRN2 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	XRN2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC,IF/ICC,ELISA
Recommended Dilution	WB 1:500-1:2000 IHC 1:50-1:200, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human XRN2
Specificity	XRN2 Antibody detects endogenous levels of XRN2
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	XRN2
Accession No.	Q9H0D6
Alternate Names	5' 3' exoribonuclease 2; DHM1 like protein; DHP protein; XRN 2;

Antibody Specification

Description	Possesses 5'→3' exoribonuclease activity (By similarity). May promote the termination of transcription by RNA polymerase II. During transcription termination, cleavage at the polyadenylation site liberates a 5' fragment which is subsequently processed to form the mature mRNA and a 3' fragment which remains attached to the elongating polymerase. The processive degradation of this 3' fragment by this protein may promote termination of transcription. Binds to RNA polymerase II (RNAP II) transcription termination R-loops formed by G-rich pause sites (PubMed:21700224).
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	108kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.