

# Immunotag™ Phospho-NF-κB p100 (Ser872) Antibody

Antibody Specification	
Catalog No.	ITA1144
Product Description	Immunotag™ Phospho-NF-κB p100 (Ser872) Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Phospho-NF-κB p100 (Ser872)
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF/ICC,ELISA
Recommended Dilution	WB 1:500-1:2000, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human NF-κB p100 around the phosphorylation site of Serine 872
Specificity	Phospho-NF-κB p100 (Ser872) Antibody detects endogenous levels of NF-κB p100 only when phosphorylated at Serine 872
Purification	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	NFKB2
Accession No.	Q00653

## Antibody Specification

Alternate Names	CVID10; DNA binding factor KBF2; H2TF1; Lymphocyte translocation chromosome 10 protein; LYT 10; NF kB2; NFKB p52/p100 subunit; Nuclear factor Kappa B subunit 2; Nuclear factor of kappa light polypeptide gene enhancer in B cells 2 (p49/p100); Nuclear factor of kappa light polypeptide gene enhancer in B cells 2; Oncogene Lys 10; p100; Transcription factor NFKB2;
Description	<p>NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. In a non-canonical activation pathway, the MAP3K14-activated CHUK/IKKA homodimer phosphorylates NFKB2/p100 associated with RelB, inducing its proteolytic processing to NFKB2/p52 and the formation of NF-kappa-B RelB-p52 complexes. The NF-kappa-B heterodimeric RelB-p52 complex is a transcriptional activator. The NF-kappa-B p52-p52 homodimer is a transcriptional repressor. NFKB2 appears to have dual functions such as cytoplasmic retention of attached NF-kappa-B proteins by p100 and generation of p52 by a cotranslational processing. The proteasome-mediated process ensures the production of both p52 and p100 and preserves their independent function. p52 binds to the kappa-B consensus sequence 5'-GGRNNYYCC-3', located in the enhancer region of genes involved in immune response and acute phase reactions. p52 and p100 are respectively the minor and major form; the processing of p100 being relatively poor. Isoform p49 is a subunit of the NF-kappa-B protein complex, which stimulates the HIV enhancer in synergy with p65. In concert with RELB, regulates the circadian clock by repressing the transcriptional activator activity of the CLOCK-ARNTL/BMAL1 heterodimer.</p>
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	100kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.