

Immunotag™ Phospho-BLNK (Tyr96) Antibody

Antibody Specification	
Catalog No.	ITA1236
Product Description	Immunotag™ Phospho-BLNK (Tyr96) Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Phospho-BLNK (Tyr96)
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC,IF/ICC,ELISA
Recommended Dilution	WB 1:500-1:2000 IHC 1:50-1:200, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human BLNK around the phosphorylation site of Tyrosine 96
Specificity	Phospho-BLNK (Tyr96) Antibody detects endogenous levels of BLNK only when phosphorylated at Tyrosine 96
Purification	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	BLNK
Accession No.	Q8WV28

Antibody Specification

Alternate Names	AGM4; B cell adapter containing SH2 domain protein; B cell adapter containing Src homology 2 domain protein; B cell adaptor containing SH2 domain; B cell linker; B cell linker protein; B cell-specific adaptor protein; B-cell activation; B-cell adapter containing a SH2 domain protein; B-cell adapter containing a Src homology 2 domain protein; B-cell linker protein; BASH; Bca; BLNK; BLNK s; BLNK_HUMAN; Cytoplasmic adapter protein; Ly 57; Ly57; Lymphocyte antigen 57; Lyw 57; Lyw57; MGC111051; SH2 domain-containing leukocyte protein, 65-KD; SLP 65; SLP-65; SLP65; Src homology [SH2] domain-containing leukocyte protein of 65 kD; Src homology 2 domain containing leukocyte protein of 65 kDa; Src homology 2 domain-containing leukocyte protein of 65 kDa;
Description	Functions as a central linker protein, downstream of the B-cell receptor (BCR), bridging the SYK kinase to a multitude of signaling pathways and regulating biological outcomes of B-cell function and development. Plays a role in the activation of ERK/EPHB2, MAP kinase p38 and JNK. Modulates AP1 activation. Important for the activation of NF-kappa-B and NFAT. Plays an important role in BCR-mediated PLCG1 and PLCG2 activation and Ca ²⁺ mobilization and is required for trafficking of the BCR to late endosomes. However, does not seem to be required for pre-BCR-mediated activation of MAP kinase and phosphatidylinositol 3 (PI3) kinase signaling. May be required for the RAC1-JNK pathway. Plays a critical role in orchestrating the pro-B cell to pre-B cell transition. May play an important role in BCR-induced B-cell apoptosis.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	50kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.