

Immunotag™ Phospho-HDAC8 (Ser39) Antibody

Antibody Specification	
Catalog No.	ITA1250
Product Description	Immunotag™ Phospho-HDAC8 (Ser39) Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Phospho-HDAC8 (Ser39)
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC,IF/ICC,ELISA
Recommended Dilution	WB 1:500-1:2000 IHC 1:50-1:200, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human HDAC8 around the phosphorylation site of Serine 39
Specificity	Phospho-HDAC8 (Ser39) Antibody detects endogenous levels of HDAC8 only when phosphorylated at Serine 39
Purification	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	HDAC8
Accession No.	Q9BY41
Alternate Names	CDA07; CDLS5; HD 8; HD8; HDAC 8; Hdac8; HDAC8_HUMAN; HDACL 1; HDACL1; Histone deacetylase 8; Histone deacetylase like 1; MRXS6; RPD 3; RPD3; WTS;

Antibody Specification

Description	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Also involved in the deacetylation of cohesin complex protein SMC3 regulating release of cohesin complexes from chromatin. May play a role in smooth muscle cell contractility.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	42kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.