

Immunotag™ Telomerase (Phospho-Ser824) Antibody

Antibody Specification	
Catalog No.	ITA1604
Product Description	Immunotag™ Telomerase (Phospho-Ser824) Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Telomerase (Phospho-Ser824)
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human Telomerase (Phospho-Ser824)
Specificity	Telomerase (Phospho-Ser824) Antibody detects endogenous levels of total Telomerase (Phospho-Ser824)
Purification	The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	TERT
Accession No.	O14746
Alternate Names	CMM9; DKCA2; DKCB4; EST2; HEST2; htert; hTRT; PFBMFT1; TCS1; Telomerase associated protein 2; Telomerase catalytic subunit; Telomerase reverse transcriptase; Telomerase-associated protein 2; Telomere Reverse Transcriptase; TERT; TERT_HUMAN; TP2; TRT;

Antibody Specification

Description	<p>Telomerase is a ribonucleoprotein enzyme essential for the replication of chromosome termini in most eukaryotes. Active in progenitor and cancer cells. Inactive, or very low activity, in normal somatic cells. Catalytic component of the telomerase holoenzyme complex whose main activity is the elongation of telomeres by acting as a reverse transcriptase that adds simple sequence repeats to chromosome ends by copying a template sequence within the RNA component of the enzyme. Catalyzes the RNA-dependent extension of 3'-chromosomal termini with the 6-nucleotide telomeric repeat unit, 5'-TTAGGG-3'. The catalytic cycle involves primer binding, primer extension and release of product once the template boundary has been reached or nascent product translocation followed by further extension. More active on substrates containing 2 or 3 telomeric repeats. Telomerase activity is regulated by a number of factors including telomerase complex-associated proteins, chaperones and polypeptide modifiers. Modulates Wnt signaling. Plays important roles in aging and antiapoptosis.</p>
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	127 kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.