Immunotag™ Synuclein-alpha Antibody

Antibody Specification	
Catalog No.	ITA1863
Product Description	Immunotag™ Synuclein-alpha Antibody
Size	100 μg, 200 μg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Synuclein-alpha
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human Synuclein-alpha
Specificity	Synuclein-alpha Antibody detects endogenous levels of total Synuclein-alpha
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt
Gene Name	SNCA
Accession No.	P37840

Antibody Specification	
Alternate Names	Alpha synuclein; Alpha-synuclein; Alpha-synuclein, isoform NACP140; alphaSYN; MGC105443; MGC110988; MGC127560; MGC64356; NACP; Non A beta component of AD amyloid; Non A4 component of amyloid precursor; Non-A beta component of AD amyloid; Non-A-beta component of alzheimers disease amyloid, precursor of; Non-A4 component of amyloid precursor; Non-A4 component of amyloid, precursor of; OTTHUMP00000218549; OTTHUMP00000218551; OTTHUMP00000218552; OTTHUMP00000218553; OTTHUMP00000218554; PARK 1; PARK 4; PARK 4; PARK 4; PARK 4; PARK 4; PARK 4; PARK 5; Parkinson disease (autosomal dominant, Lewy body) 4; Parkinson disease familial 1; SNCA; Snca synuclein, alpha (non A4 component of amyloid precursor); SYN; Synuclein alpha; Synuclein alpha 140; Synuclein, alpha (non A4 component of amyloid precursor); SYUA_HUMAN;
Description	May be involved in the regulation of dopamine release and transport. Induces fibrillization of microtubule-associated protein tau. Reduces neuronal responsiveness to various apoptotic stimuli, leading to a decreased caspase-3 activation.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	18kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.

www.gbiosciences.com

© 2018 Geno Technology Inc., USA. All Rights Reserved.