

Immunotag™ TAAR1 Antibody

| Antibody Specification | |
|------------------------|--|
| Catalog No. | ITA2558 |
| Product Description | Immunotag™ TAAR1 Antibody |
| Size | 100 µg, 200 µg |
| Conjugation | HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647 |
| IMPORTANT NOTE | This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return. |
| Target Protein | TAAR1 |
| Clonality | Polyclonal |
| Storage/Stability | -20°C/1 year |
| Application | WB,IHC,ELISA |
| Recommended Dilution | WB 1:1000-3000 IHC 1:200 |
| Concentration | 1 mg/ml |
| Reactive Species | Human,Mouse,Rat |
| Host Species | Rabbit |
| Immunogen | A synthesized peptide derived from human TAAR1 |
| Specificity | TAAR1 Antibody detects endogenous levels of total TAAR1 |
| Purification | The antiserum was purified by peptide affinity chromatography. |
| Form | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt |
| Gene Name | TAAR1 |
| Accession No. | Q96RJ0 |
| Alternate Names | Taar1; TAAR1_HUMAN; TaR-1; Trace amine associated receptor 1; Trace amine receptor 1; Trace amine-associated receptor 1; Trar1; |

Antibody Specification

| | |
|---------------------------|--|
| Description | Receptor for trace amines, including beta-phenylethylamine (b-PEA), p-tyramine (p-TYR), octopamine and tryptamine, with highest affinity for b-PEA and p-TYR. Unresponsive to classical biogenic amines, such as epinephrine and histamine and only partially activated by dopamine and serotonin. Trace amines are biogenic amines present in very low levels in mammalian tissues. Although some trace amines have clearly defined roles as neurotransmitters in invertebrates, the extent to which they function as true neurotransmitters in vertebrates has remained speculative. Trace amines are likely to be involved in a variety of physiological functions that have yet to be fully understood. The signal transduced by this receptor is mediated by the G(s)-class of G-proteins which activate adenylate cyclase. |
| Cell Pathway/ Category | Primary Polyclonal Antibody |
| Protein MW | 39 kDa |
| Usage | For Research Use Only! Not for diagnostic or therapeutic procedures. |