

Immunotag™ TET2 Antibody

Antibody Specification	
Catalog No.	ITA2743
Product Description	Immunotag™ TET2 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	TET2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC
Recommended Dilution	WB 1:500-1:2000, IHC 1:50-1:200
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human TET2.
Specificity	TET2 antibody detects endogenous levels of TET2.
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	TET2
Accession No.	Q6N021
Alternate Names	FLJ20032; KIAA1546; MDS; Methylcytosine dioxygenase TET2; Nbla00191; Probable methylcytosine dioxygenase TET2; Protein Ayu17 449; Tet 2; Tet methylcytosine dioxygenase 2; Tet oncogene 2; Tet oncogene family member 2; TET2; TET2_HUMAN;

Antibody Specification

Description	Dioxygenase that catalyzes the conversion of the modified genomic base 5-methylcytosine (5mC) into 5-hydroxymethylcytosine (5hmC) and plays a key role in active DNA demethylation. Has a preference for 5-hydroxymethylcytosine in CpG motifs. Also mediates subsequent conversion of 5hmC into 5-formylcytosine (5fC), and conversion of 5fC to 5-carboxylcytosine (5caC). Conversion of 5mC into 5hmC, 5fC and 5caC probably constitutes the first step in cytosine demethylation. Methylation at the C5 position of cytosine bases is an epigenetic modification of the mammalian genome which plays an important role in transcriptional regulation. In addition to its role in DNA demethylation, also involved in the recruitment of the O-GlcNAc transferase OGT to CpG-rich transcription start sites of active genes, thereby promoting histone H2B GlcNAcylation by OGT.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	130 kDa,133 kDa,224 kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.