

# Immunotag™ RBM4 Antibody

Antibody Specification	
Catalog No.	ITA2879
Product Description	Immunotag™ RBM4 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	RBM4
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC
Recommended Dilution	WB 1:500-1:2000, IHC 1:50-1:200
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human RBM4.
Specificity	RBM4 antibody detects endogenous levels of RBM4.
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	RBM4
Accession No.	Q9BWF3
Alternate Names	dkfzp547k0918; hLark; lark; Lark homolog; lark homologue; mgc75138; RBM4; rbm4 lark; RBM4_HUMAN; RNA binding motif 4; RNA binding motif 4a; RNA-binding motif protein 4; RNA-binding motif protein 4a; RNA-binding protein 4;

## Antibody Specification

Description	RNA-binding factor involved in multiple aspects of cellular processes like alternative splicing of pre-mRNA and translation regulation. Modulates alternative 5'-splice site and exon selection. Acts as a muscle cell differentiation-promoting factor. Activates exon skipping of the PTB pre-mRNA during muscle cell differentiation. Antagonizes the activity of the splicing factor PTBP1 to modulate muscle cell-specific exon selection of alpha tropomyosin. Binds to intronic pyrimidine-rich sequence of the TPM1 and MAPT pre-mRNAs. Required for the translational activation of PER1 mRNA in response to circadian clock. Binds directly to the 3'-UTR of the PER1 mRNA. Exerts a suppressive activity on Cap-dependent translation via binding to CU-rich responsive elements within the 3'UTR of mRNAs, a process increased under stress conditions or during myocytes differentiation. Recruits EIF4A1 to stimulate IRES-dependent translation initiation in response to cellular stress. Associates to internal ribosome entry segment (IRES) in target mRNA species under stress conditions. Plays a role for miRNA-guided RNA cleavage and translation suppression by promoting association of AGO2-containing miRNPs with their cognate target mRNAs. Associates with miRNAs during muscle cell differentiation. Binds preferentially to 5'-CGCGCG[GCA]-3' motif in vitro.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	40 kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.