

Immunotag™ SH2D1B Antibody

Antibody Specification	
Catalog No.	ITA2972
Product Description	Immunotag™ SH2D1B Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	SH2D1B
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC
Recommended Dilution	WB 1:500-1:2000, IHC 1:50-1:200
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human SH2D1B.
Specificity	SH2D1B antibody detects endogenous levels of SH2D1B.
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	SH2D1B
Accession No.	O14796
Alternate Names	EAT 2; EAT2; EWS/FLI1 activated transcript 2; SH2 domain containing 1B; SH2 domain containing molecule EAT2; SH2 domain containing protein 1B;

Antibody Specification

Description	Cytoplasmic adapter regulating receptors of the signaling lymphocytic activation molecule (SLAM) family such as CD84, SLAMF1, LY9 and CD244 (PubMed:11689425). In SLAM signaling seems to cooperate with SH2D1A/SAP. Plays a role in regulation of effector functions of natural killer (NK) cells by controlling signal transduction through CD244/2B4 without effecting its tyrosine phosphorylation; downstream signaling involves PLCG1 and ERK activation (PubMed:24687958). Activation of SLAMF7-mediated NK cell function does not effect receptor tyrosine phosphorylation but distal signaling (By similarity). In the context of NK cell-mediated cytotoxicity does not enhance conjugate formation with target cells but stimulates polarization of the microtubule-organizing center and cytotoxic granules toward the NK cell synapse (PubMed:24687958). Negatively regulates CD40-induced cytokine production in dendritic cells downstream of SLAM family receptors probably by inducing activation of the PI3K pathway to inhibit p38 MAPK and JNK activation (By similarity).
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	15 kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.