

## Immunotag™ PIK3CA Antibody

Antibody Specification	
Catalog No.	ITA3565
Product Description	Immunotag™ PIK3CA Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PIK3CA
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	A synthesized peptide
Specificity	PIK3CA antibody detects endogenous levels of total PIK3CA
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Gene Name	PIK3CA
Accession No.	P42336

## Antibody Specification

Alternate Names	5-bisphosphate 3-kinase 110 kDa catalytic subunit alpha; 5-bisphosphate 3-kinase catalytic subunit alpha isoform; caPI3K; CLOVE; CWS5; MCAP; MCM; MCMTc; MGC142161; MGC142163; p110 alpha; p110alpha; Phosphatidylinositol 3 kinase catalytic alpha polypeptide; Phosphatidylinositol 3 kinase catalytic 110 KD alpha; Phosphatidylinositol 4 5 bisphosphate 3 kinase catalytic subunit alpha; Phosphatidylinositol 4,5 bisphosphate 3 kinase catalytic subunit alpha isoform; Phosphatidylinositol 4,5 bisphosphate 3 kinase 110 kDa catalytic subunit alpha; Phosphatidylinositol-4; Phosphoinositide 3 kinase catalytic alpha polypeptide; PI3 kinase p110 subunit alpha; PI3-kinase subunit alpha; PI3K; PI3K-alpha; PI3KC A; PIK3C A; Pik3ca; PK3CA; PK3CA_HUMAN; PtdIns 3 kinase p110; PtdIns-3-kinase subunit alpha; PtdIns-3-kinase subunit p110-alpha; Serine/threonine protein kinase PIK3CA;
Description	Phosphoinositide-3-kinase (PI3K) that phosphorylates PtdIns (Phosphatidylinositol), PtdIns4P (Phosphatidylinositol 4-phosphate) and PtdIns(4,5)P2 (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3). PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Participates in cellular signaling in response to various growth factors. Involved in the activation of AKT1 upon stimulation by receptor tyrosine kinases ligands such as EGF, insulin, IGF1, VEGFA and PDGF. Involved in signaling via insulin-receptor substrate (IRS) proteins. Essential in endothelial cell migration during vascular development through VEGFA signaling, possibly by regulating RhoA activity. Required for lymphatic vasculature development, possibly by binding to RAS and by activation by EGF and FGF2, but not by PDGF. Regulates invadopodia formation through the PDK1-AKT1 pathway. Participates in cardiomyogenesis in embryonic stem cells through a AKT1 pathway. Participates in vasculogenesis in embryonic stem cells through PDK1 and protein kinase C pathway. Also has serine-protein kinase activity: phosphorylates PIK3R1 (p85alpha regulatory subunit), EIF4EBP1 and HRAS. Plays a role in the positive regulation of phagocytosis and pinocytosis (By similarity).
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	110 kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.