

Immunotag™ GPR40 Antibody

Antibody Specification	
Catalog No.	ITA3684
Product Description	Immunotag™ GPR40 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	GPR40
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide
Specificity	GPR40 antibody detects endogenous levels of total GPR40
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Gene Name	FFAR1
Accession No.	O14842
Alternate Names	FFA1R; FFAR 1; Ffar1; FFAR1_HUMAN; Free fatty acid receptor 1; G protein coupled receptor 40; G-protein coupled receptor 40; GPCR40; GPR 40; GPR40;

Antibody Specification

Description	G-protein coupled receptor for medium and long chain saturated and unsaturated fatty acids that plays an important role in glucose homeostasis. Fatty acid binding increases glucose-stimulated insulin secretion, and may also enhance the secretion of glucagon-like peptide 1 (GLP-1). May also play a role in bone homeostasis; receptor signaling activates pathways that inhibit osteoclast differentiation (By similarity). Ligand binding leads to a conformation change that triggers signaling via G-proteins that activate phospholipase C, leading to an increase of the intracellular calcium concentration. Seems to act through a G(q) and G(i)-mediated pathway.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	28 kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.