

## Immunotag™ PSMD2 Antibody

Antibody Specification	
Catalog No.	ITA4170
Product Description	Immunotag™ PSMD2 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PSMD2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF/ICC,ELISA
Recommended Dilution	WB 1:500~1:1000, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide
Specificity	PSMD2 Antibody detects endogenous levels of total PSMD2
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	PSMD2
Accession No.	Q13200
Alternate Names	26S proteasome non-ATPase regulatory subunit 2; 26S proteasome regulatory subunit RPN1; 26S proteasome regulatory subunit S2; 26S proteasome subunit p97; 55.11 protein; MGC14274; P97; Proteasome (prosome macropain) 26S subunit non ATPase 2; Proteasome 26S subunit, non-ATPase, 2; Proteasome 26S, subunit 2; Protein 55.11; Psmd2; PSMD2_HUMAN; Rpn1; S2; TNFR associated protein 2; TRAP2; Tumor necrosis factor receptor associated protein 2; Tumor necrosis factor type 1 receptor-associated protein 2;

## Antibody Specification

Description	Component of the 26S proteasome, a multiprotein complex involved in the ATP-dependent degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins, which could impair cellular functions, and by removing proteins whose functions are no longer required. Therefore, the proteasome participates in numerous cellular processes, including cell cycle progression, apoptosis, or DNA damage repair.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	100 KD
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.