

Immunotag™ SIAH1 Antibody

Antibody Specification	
Catalog No.	ITA4227
Product Description	Immunotag™ SIAH1 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	SIAH1
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF/ICC,ELISA
Recommended Dilution	WB 1:500~1:1000, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide
Specificity	SIAH1 Antibody detects endogenous levels of total SIAH1
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	SIAH1
Accession No.	Q8IUQ4/O43255

Antibody Specification

Alternate Names	E3 ubiquitin-protein ligase SIAH1; FLJ08065; hSIAH1; HUMSIAH; Seven in absentia homolog 1 (Drosophila); Seven in absentia homolog 1; Siah 1; Siah 1a; Siah E3 ubiquitin protein ligase 1; Siah-1; Siah-1a; Siah1; SIAH1_HUMAN; SIAH1A; Ubiquitin ligase SIAH1; E3 ubiquitin protein ligase SIAH2; E3 ubiquitin-protein ligase SIAH1; hSIAH1; hSiah2; HUMSIAH; Seven in absentia homolog 1 (Drosophila); Seven in absentia homolog 1; Seven in Absentia Homolog 2; SIAH 1; Siah 1a; Siah 2; Siah-1; Siah-1a; Siah1; SIAH1_HUMAN; Ubiquitin ligase SIAH1; Ubiquitin Ligase Siah2;
Description	E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent proteasomal degradation of target proteins. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Mediates E3 ubiquitin ligase activity either through direct binding to substrates or by functioning as the essential RING domain subunit of larger E3 complexes. Triggers the ubiquitin-mediated degradation of many substrates, including proteins involved in transcription regulation (ELL2, MYB, POU2AF1, PML and RBBP8), a cell surface receptor (DCC), the cell-surface receptor-type tyrosine kinase FLT3, the cytoplasmic signal transduction molecules (KLF10/TIEG1 and NUMB), an antiapoptotic protein (BAG1), a microtubule motor protein (KIF22), a protein involved in synaptic vesicle function in neurons (SYP), a structural protein (CTNNB1) and SNCAIP. Confers constitutive instability to HIPK2 through proteasomal degradation. It is thereby involved in many cellular processes such as apoptosis, tumor suppression, cell cycle, axon guidance, transcription regulation, spermatogenesis and TNF-alpha signaling. Has some overlapping function with SIAH2. Induces apoptosis in cooperation with PEG3. Upon nitric oxid (NO) generation that follows apoptotic stimulation, interacts with S-nitrosylated GAPDH, mediating the translocation of GAPDH to the nucleus. GAPDH acts as a stabilizer of SIAH1, facilitating the degradation of nuclear proteins.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	34 KD
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.