

Immunotag™ IKZF3 Antibody

Antibody Specification	
Catalog No.	ITA4356
Product Description	Immunotag™ IKZF3 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	IKZF3
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF/ICC,ELISA
Recommended Dilution	WB 1:500~1:1000, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	A synthesized peptide
Specificity	IKZF3 Antibody detects endogenous levels of total IKZF3
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	IKZF3
Accession No.	Q9UKT9
Alternate Names	AIO; Aiolos; IKAROS family zinc finger 3 (Aiolos); IKAROS family zinc finger 3; Ikaros family zinc finger protein 3; IKZF 3; IKZF3; IKZF3_HUMAN; zinc finger DNA binding protein Aiolos; Zinc finger protein Aiolos; Zinc finger protein subfamily 1A 3 (Aiolos); Zinc finger protein subfamily 1A 3; Zinc finger protein subfamily 1A, member 3; ZNFN1A3;

Antibody Specification

Description	Transcription factor that plays an important role in the regulation of lymphocyte differentiation. Plays an essential role in regulation of B-cell differentiation, proliferation and maturation to an effector state. Involved in regulating BCL2 expression and controlling apoptosis in T-cells in an IL2-dependent manner.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	62 KD
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.