

Immunotag™ TAF3 Antibody

Antibody Specification	
Catalog No.	ITA5453
Product Description	Immunotag™ TAF3 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	TAF3
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500~1:1000
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide
Specificity	TAF3 Antibody detects endogenous levels of total TAF3
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	TAF3
Accession No.	Q5VWG9
Alternate Names	140 kDa TATA box-binding protein-associated factor; Bip2 protein; RNA polymerase II transcription factor TAFII140; TAF(II)140; TAF140; TAF3; TAF3 RNA polymerase II TATA box binding protein (TBP)-associated factor; TAF3_HUMAN; TAFII-140; TAFII140; TAFII155 protein; TATA box binding protein associated factor 3; TBP-associated factor 3; Transcription initiation factor TFIID 140 kDa subunit; Transcription initiation factor TFIID subunit 3;

Antibody Specification

Description	Transcription factor TFIID is one of the general factors required for accurate and regulated initiation by RNA polymerase II. TFIID is a multimeric protein complex that plays a central role in mediating promoter responses to various activators and repressors. Required in complex with TBPL2 for the differentiation of myoblasts into myocytes. The complex replaces TFIID at specific promoters at an early stage in the differentiation process.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	100 KD
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.