

## Immunotag™ UBE2T Antibody

| Antibody Specification |   |
|------------------------|---|
| Catalog No.            | ITA5479   |
| Product Description    | Immunotag™ UBE2T Antibody   |
| Size                   | 100 µg, 200 µg  |
| Conjugation            | HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647   |
| IMPORTANT NOTE         | This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.  |
| Target Protein         | UBE2T   |
| Clonality              | Polyclonal  |
| Storage/Stability      | -20°C/1 year  |
| Application            | WB,IF/ICC,ELISA   |
| Recommended Dilution   | WB 1:500~1:1000, IF/ICC 1:100-1:500   |
| Concentration          | 1 mg/ml   |
| Reactive Species       | Human,Mouse   |
| Host Species           | Rabbit  |
| Immunogen              | A synthesized peptide   |
| Specificity            | UBE2T Antibody detects endogenous levels of total UBE2T   |
| Purification           | The antiserum was purified by peptide affinity chromatography.  |
| Form                   | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt   |
| Gene Name              | UBE2T   |
| Accession No.          | Q9NPD8  |
| Alternate Names        | Cell proliferation inducing gene 50 protein; Cell proliferation-inducing gene 50 protein; HSPC150; HSPC150 protein similar to ubiquitin conjugating enzyme; PIG50; Ube2t; UBE2T_HUMAN; Ubiquitin carrier protein T; Ubiquitin conjugating enzyme; ubiquitin conjugating enzyme E2 T; Ubiquitin conjugating enzyme E2T; Ubiquitin protein ligase T; Ubiquitin-conjugating enzyme E2 T; Ubiquitin-protein ligase T; |

## Antibody Specification

|                           |  |
|---------------------------|--|
| Description               | Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. Catalyzes monoubiquitination. Involved in mitomycin-C (MMC)-induced DNA repair. Acts as a specific E2 ubiquitin-conjugating enzyme for the Fanconi anemia complex by associating with E3 ubiquitin-protein ligase FANCL and catalyzing monoubiquitination of FANCD2, a key step in the DNA damage pathway (PubMed:16916645, PubMed:17938197, PubMed:19111657, PubMed:19589784, PubMed:28437106). Also mediates monoubiquitination of FANCL and FANCI (PubMed:16916645, PubMed:17938197, PubMed:19111657, PubMed:19589784). May contribute to ubiquitination and degradation of BRCA1 (PubMed:19887602). In vitro able to promote polyubiquitination using all 7 ubiquitin Lys residues, but may prefer 'Lys-11', 'Lys-27', 'Lys-48' and 'Lys-63'-linked polyubiquitination (PubMed:20061386). |
| Cell Pathway/<br>Category | Primary Polyclonal Antibody  |
| Protein MW                | 22 KD  |
| Usage                     | For Research Use Only! Not for diagnostic or therapeutic procedures.   |