

Immunotag™ NDFIP1 Antibody

Antibody Specification	
Catalog No.	ITA5559
Product Description	Immunotag™ NDFIP1 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	NDFIP1
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF/ICC,ELISA
Recommended Dilution	WB 1:500~1:1000, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide
Specificity	NDFIP1 Antibody detects endogenous levels of total NDFIP1
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	NDFIP1
Accession No.	Q9BT67

Antibody Specification

Alternate Names	0610010M22Rik; Breast cancer associated protein SGA 1M; Breast cancer-associated protein SGA-1M; MGC10924; N4WBP5; Ndfip1; Nedd4 family interacting protein 1; NEDD4 family-interacting protein 1; NEDD4 WW domain binding protein 5; NEDD4 WW domain-binding protein 5; NFIP1_HUMAN; Putative MAPK activating protein PM13; Putative MAPK-activating protein PM13; Putative NF kappa B activating protein 164; Putative NF-kappa-B-activating protein 164; Putative NFKB and MAPK activating protein; Putative NFKB and MAPK-activating protein;
Description	<p>Activates HECT domain-containing E3 ubiquitin-protein ligases, including NEDD4 and ITCH, and consequently modulates the stability of their targets. As a result, controls many cellular processes. Prevents chronic T-helper cell-mediated inflammation by activating ITCH and thus controlling JUNB degradation (By similarity). Promotes pancreatic beta cell death through degradation of JUNB and inhibition of the unfolded protein response, leading to reduction of insulin secretion (PubMed:26319551). Restricts the production of proinflammatory cytokines in effector Th17 T-cells by promoting ITCH-mediated ubiquitination and degradation of RORC (By similarity). Together with NDFIP2, limits the cytokine signaling and expansion of effector Th2 T-cells by promoting degradation of JAK1, probably by ITCH- and NEDD4L-mediated ubiquitination (By similarity). Regulates peripheral T-cell tolerance to self and foreign antigens, forcing the exit of naive CD4+ T-cells from the cell cycle before they become effector T-cells (By similarity). Negatively regulates RLR-mediated antiviral response by promoting SMURF1-mediated ubiquitination and subsequent degradation of MAVS (PubMed:23087404). Negatively regulates KCNH2 potassium channel activity by decreasing its cell-surface expression and interfering with channel maturation through recruitment of NEDD4L to the Golgi apparatus where it mediates KCNH2 degradation (PubMed:26363003). In cortical neurons, mediates the ubiquitination of the divalent metal transporter SLC11A2/DMT1 by NEDD4L, leading to its down-regulation and protection of the cells from cobalt and iron toxicity (PubMed:19706893). Important for normal development of dendrites and dendritic spines in cortex (By similarity). Enhances the ubiquitination of BRAT1 mediated by: NEDD4, NEDD4L and ITCH and is required for the nuclear localization of ubiquitinated BRAT1 (PubMed:25631046). Enhances the ITCH-mediated ubiquitination of MAP3K7 by recruiting E2 ubiquitin-conjugating enzyme UBE2L3 to ITCH (By similarity). Modulates EGFR signaling through multiple pathways. In particular, may regulate the ratio of AKT1-to-MAPK8 signaling in response to EGF, acting on AKT1 probably through PTEN destabilization and on MAPK8 through ITCH-dependent MAP2K4 inactivation. As a result, may control cell growth rate (PubMed:20534535). Inhibits cell proliferation by promoting PTEN nuclear localization and changing its signaling specificity (PubMed:25801959).</p>
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	25 KD
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.