## Immunotag<sup>™</sup> CDC20 Antibody

## **Antibody Specification**

Catalog No.	ITA6445
Product Description	Immunotag™ CDC20 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	CDC20
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC,IF/ICC,ELISA
Recommended Dilution	WB 1:500-1:2000 IHC 1:50-1:200, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human CDC20
Specificity	CDC20 Antibody detects endogenous levels of total CDC20
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	CDC20
Accession No.	Q12834
Alternate Names	bA276H19.3; Cdc 20; CDC20; CDC20 cell division cycle 20 homolog; CDC20_HUMAN; CDC20A; Cell division cycle 20; Cell division cycle 20 homolog (S. cerevisiae); Cell division cycle 20 homolog; Cell division cycle protein 20 homolog; fizzy; MGC102824; p55CDC;

## **Antibody Specification**

Description	Required for full ubiquitin ligase activity of the anaphase promoting complex/cyclosome (APC/C) and may confer substrate specificity upon the complex. Is regulated by MAD2L1: in metaphase the MAD2L1-CDC20-APC/C ternary complex is inactive and in anaphase the CDC20-APC/C binary complex is active in degrading substrates. The CDC20-APC/C complex positively regulates the formation of synaptic vesicle clustering at active zone to the presynaptic membrane in postmitotic neurons. CDC20-APC/C-induced degradation of NEUROD2 induces presynaptic differentiation.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	55kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.

www.gbiosciences.com

© 2018 Geno Technology Inc., USA. All Rights Reserved.