

Immunotag™ MAX Antibody

Antibody Specification	
Catalog No.	ITA6957
Product Description	Immunotag™ MAX Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	MAX
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC,ELISA
Recommended Dilution	WB 1:500-1:2000 IHC 1:50-1:200
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human MAX
Specificity	MAX Antibody detects endogenous levels of total MAX
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	MAX
Accession No.	P61244
Alternate Names	bHLHd4; bHLHd5; bHLHd6; bHLHd7; bHLHd8; Helix loop helix zipper protein; MAX protein; MGC10775; MGC11225; MGC18164; MGC34679; MGC36767; MYC associated factor X; Myc binding novel HLH/LZ protein; Orf 1; Orf1; Protein max;

Antibody Specification

Description	Transcription regulator. Forms a sequence-specific DNA-binding protein complex with MYC or MAD which recognizes the core sequence 5'-CAC[GA]TG-3'. The MYC:MAX complex is a transcriptional activator, whereas the MAD:MAX complex is a repressor. May repress transcription via the recruitment of a chromatin remodeling complex containing H3 'Lys-9' histone methyltransferase activity. Represses MYC transcriptional activity from E-box elements.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	18kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.