

# Immunotag™ RPA2 Antibody

| Antibody Specification |  |
|------------------------|--|
| Catalog No.            | ITA6989  |
| Product Description    | Immunotag™ RPA2 Antibody   |
| Size                   | 100 µg, 200 µg   |
| Conjugation            | HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647                                      |
| IMPORTANT NOTE         | This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return. |
| Target Protein         | RPA2   |
| Clonality              | Polyclonal   |
| Storage/Stability      | -20°C/1 year   |
| Application            | WB,IHC,IF/ICC,ELISA  |
| Recommended Dilution   | WB 1:500-1:2000 IHC 1:50-1:200, IF/ICC 1:100-1:500   |
| Concentration          | 1 mg/ml  |
| Reactive Species       | Human,Mouse,Rat  |
| Host Species           | Rabbit   |
| Immunogen              | A synthesized peptide derived from human RPA2  |
| Specificity            | RPA2 Antibody detects endogenous levels of total RPA2  |
| Purification           | The antiserum was purified by peptide affinity chromatography.   |
| Form                   | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt        |
| Gene Name              | RPA2   |
| Accession No.          | P15927   |

## Antibody Specification

|                           |  |
|---------------------------|--|
| Alternate Names           | 60S acidic ribosomal protein P1; AA409079; AI325195; AU020965; ik:tdsubc_2g1; M(2)21C; MGC137236; OTTHUMP00000004008; p32; p34; RCJMB04_6d17 replication protein A2, 32kDa; REPA2; Replication factor A protein 2; Replication protein A 32 kDa subunit; Replication protein A 32kDa subunit; Replication protein A 34 kDa subunit; Replication protein A; Replication Protein A2 (32kDa); Replication protein A2; Replication protein A2, 32kDa; RF-A protein 2; Rf-A2; RFA; RFA2_HUMAN; RP-A p32; RP-A p34; RP21C; RPA 2; RPA 32; RPA; Rpa2; RPA32; RPA34; RpLP1; RpP2; xx:tdsubc_2g1; zgc:109822;   |
| Description               | As part of the heterotrimeric replication protein A complex (RPA/RP-A), binds and stabilizes single-stranded DNA intermediates, that form during DNA replication or upon DNA stress. It prevents their reannealing and in parallel, recruits and activates different proteins and complexes involved in DNA metabolism. Thereby, it plays an essential role both in DNA replication and the cellular response to DNA damage. In the cellular response to DNA damage, the RPA complex controls DNA repair and DNA damage checkpoint activation. Through recruitment of ATRIP activates the ATR kinase a master regulator of the DNA damage response. It is required for the recruitment of the DNA double-strand break repair factors RAD51 and RAD52 to chromatin in response to DNA damage. Also recruits to sites of DNA damage proteins like XPA and XPG that are involved in nucleotide excision repair and is required for this mechanism of DNA repair. Plays also a role in base excision repair (BER) probably through interaction with UNG. Also recruits SMARCAL1/HARP, which is involved in replication fork restart, to sites of DNA damage. May also play a role in telomere maintenance. |
| Cell Pathway/<br>Category | Primary Polyclonal Antibody  |
| Protein MW                | 29kDa  |
| Usage                     | For Research Use Only! Not for diagnostic or therapeutic procedures.   |