

Immunotag™ RBPJ Antibody

Antibody Specification	
Catalog No.	ITA7513
Product Description	Immunotag™ RBPJ Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	RBPJ
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC,ELISA
Recommended Dilution	WB 1:500-1:2000 IHC 1:50-1:200
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human RBPJ
Specificity	RBPJ Antibody detects endogenous levels of total RBPJ
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	RBPJ
Accession No.	Q06330
Alternate Names	AI843960; AOS3; CBF 1; CBF-1; CBF1; csl; IGKJRB; IGKJRB1; J kappa recombination signal binding protein; J kappa-recombination signal-binding protein; KBF2; NY REN 30 antigen; RBP J; RBP J kappa; RBP JK; RBP-J; RBP-J kappa; RBP-JK; Rbpj; RBPJK; RBPSUH; recombination signal binding protein for immunoglobulin kappa J region; Recombining binding protein suppressor of hairless; Renal carcinoma antigen NY-REN-30; SUH; SUH_HUMAN;

## Antibody Specification

Description	Transcriptional regulator that plays a central role in Notch signaling, a signaling pathway involved in cell-cell communication that regulates a broad spectrum of cell-fate determinations. Acts as a transcriptional repressor when it is not associated with Notch proteins. When associated with some NICD product of Notch proteins (Notch intracellular domain), it acts as a transcriptional activator that activates transcription of Notch target genes. Probably represses or activates transcription via the recruitment of chromatin remodeling complexes containing histone deacetylase or histone acetylase proteins, respectively. Specifically binds to the immunoglobulin kappa-type J segment recombination signal sequence. Binds specifically to methylated DNA (PubMed:21991380). Binds to the oxygen responsive element of COX4I2 and activates its transcription under hypoxia conditions (4% oxygen) (PubMed:23303788). Negatively regulates the phagocyte oxidative burst in response to bacterial infection by repressing transcription of NADPH oxidase subunits (By similarity).
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	55kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.