

Immunotag™ PRMT7 Antibody

Antibody Specification	
Catalog No.	ITA7756
Product Description	Immunotag™ PRMT7 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PRMT7
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF/ICC,ELISA
Recommended Dilution	WB 1:1000-3000, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human PRMT7
Specificity	PRMT7 Antibody detects endogenous levels of total PRMT7
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	PRMT7
Accession No.	Q9NVM4
Alternate Names	[Myelin basic protein]-arginine N-methyltransferase PRMT7; ANM7_HUMAN; FLJ10640; Histone-arginine N-methyltransferase PRMT7; KIAA1933; Myelin basic protein arginine N methyltransferase; OTTHUMP00000174863; PRMT7; Protein arginine methyltransferase 7; Protein arginine N-methyltransferase 7;

Antibody Specification

Description	Arginine methyltransferase that can both catalyze the formation of omega-N monomethylarginine (MMA) and symmetrical dimethylarginine (sDMA), with a preference for the formation of MMA. Specifically mediates the symmetrical dimethylation of arginine residues in the small nuclear ribonucleoproteins Sm D1 (SNRPD1) and Sm D3 (SNRPD3); such methylation being required for the assembly and biogenesis of snRNP core particles. Specifically mediates the symmetric dimethylation of histone H4 'Arg-3' to form H4R3me2s. Plays a role in gene imprinting by being recruited by CTCFL at the H19 imprinted control region (ICR) and methylating histone H4 to form H4R3me2s, possibly leading to recruit DNA methyltransferases at these sites. May also play a role in embryonic stem cell (ESC) pluripotency. Also able to mediate the arginine methylation of histone H2A and myelin basic protein (MBP) in vitro; the relevance of such results is however unclear in vivo.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	78 kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.