

## Immunotag™ NR2C2 Antibody

| Antibody Specification |                                                                                                                                                                                                                                                                          |
|------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Catalog No.            | ITA8029                                                                                                                                                                                                                                                                  |
| Product Description    | Immunotag™ NR2C2 Antibody                                                                                                                                                                                                                                                |
| Size                   | 100 µg, 200 µg                                                                                                                                                                                                                                                           |
| Conjugation            | HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647                                                                                                                                            |
| IMPORTANT NOTE         | This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.                                                                                                       |
| Target Protein         | NR2C2                                                                                                                                                                                                                                                                    |
| Clonality              | Polyclonal                                                                                                                                                                                                                                                               |
| Storage/Stability      | -20°C/1 year                                                                                                                                                                                                                                                             |
| Application            | WB,ELISA                                                                                                                                                                                                                                                                 |
| Recommended Dilution   | WB 1:1000-3000                                                                                                                                                                                                                                                           |
| Concentration          | 1 mg/ml                                                                                                                                                                                                                                                                  |
| Reactive Species       | Human                                                                                                                                                                                                                                                                    |
| Host Species           | Rabbit                                                                                                                                                                                                                                                                   |
| Immunogen              | A synthesized peptide derived from human NR2C2                                                                                                                                                                                                                           |
| Specificity            | NR2C2 Antibody detects endogenous levels of total NR2C2                                                                                                                                                                                                                  |
| Purification           | The antiserum was purified by peptide affinity chromatography.                                                                                                                                                                                                           |
| Form                   | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt                                                                                                              |
| Gene Name              | NR2C2                                                                                                                                                                                                                                                                    |
| Accession No.          | P49116                                                                                                                                                                                                                                                                   |
| Alternate Names        | hTAK1; Nr2c2; NR2C2_HUMAN; Nuclear hormone receptor TR4; Nuclear receptor subfamily 2 group C member 2; Orphan nuclear receptor TAK1; Orphan nuclear receptor TR4; TAK1; Testicular nuclear receptor 4; Testicular receptor 4; TR2R1; TR4; TR4 nuclear hormone receptor; |

## Antibody Specification

|                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|---------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Description               | Orphan nuclear receptor that can act as a repressor or activator of transcription. An important repressor of nuclear receptor signaling pathways such as retinoic acid receptor, retinoid X, vitamin D3 receptor, thyroid hormone receptor and estrogen receptor pathways. May regulate gene expression during the late phase of spermatogenesis. Together with NR2C1, forms the core of the DRED (direct repeat erythroid-definitive) complex that represses embryonic and fetal globin transcription including that of GATA1. Binds to hormone response elements (HREs) consisting of two 5'-AGGTCA-3' half site direct repeat consensus sequences. Plays a fundamental role in early embryonic development and embryonic stem cells. Required for normal spermatogenesis and cerebellum development. Appears to be important for neurodevelopmentally regulated behavior (By similarity). Activates transcriptional activity of LHCG. Antagonist of PPARA-mediated transactivation. |
| Cell Pathway/<br>Category | Primary Polyclonal Antibody                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Protein MW                | 67 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Usage                     | For Research Use Only! Not for diagnostic or therapeutic procedures.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |