Immunotag[™] IFITM3 Antibody

Antibody Specification

Catalog No.	ITA8095
Product Description	Immunotag™ IFITM3 Antibody
Size	100 μg, 200 μg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	IFITM3
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IF/ICC,ELISA
Recommended Dilution	WB 1:1000-3000, IF/ICC 1:100-1:500
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human IFITM3
Specificity	IFITM3 Antibody detects endogenous levels of total IFITM3
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	IFITM3
Accession No.	Q01628
Alternate Names	1 8U; Fragilis; IFITM3; IFM3_HUMAN; Interferon induced transmembrane protein 3 (1 8U); Interferon Induced Transmembrane Protein 3; Interferon inducible; Interferon inducible protein 1 8U; Interferon Inducible Protein 15; Interferon Inducible Protein Homolog; Interferon-induced transmembrane protein 3; Interferon-inducible protein 1-8U; IP15;

Antibody Specification

Description	IFN-induced antiviral protein which disrupts intracellular cholesterol homeostasis. Inhibits the entry of viruses to the host cell cytoplasm by preventing viral fusion with cholesterol depleted endosomes. May inactivate new enveloped viruses which buds out of the infected cell, by letting them go out with a cholesterol depleted membrane. Active against multiple viruses, including influenza A virus, SARS coronavirus (SARS-CoV), Marburg virus (MARV) and Ebola virus (EBOV), Dengue virus (DNV), West Nile virus (WNV), human immunodeficiency virus type 1 (HIV-1) and vesicular stomatitis virus (VSV). Can inhibit: influenza virus hemagglutinin protein-mediated viral entry, MARV and EBOV GP1,2-mediated viral entry, SARS-CoV S protein-mediated viral entry and VSV G protein-mediated viral entry. Plays a critical role in the structural stability and function of vacuolar ATPase (v-ATPase). Establishes physical contact with the v-ATPase of endosomes which is critical for proper clathrin localization and is also required for the function of the v-ATPase to lower the pH in phagocytic endosomes thus establishing an antiviral state.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	15 kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.

www.gbiosciences.com

 $\ensuremath{\mathbb{C}}$ 2018 Geno Technology Inc., USA. All Rights Reserved.