

Immunotag™ RPC7 Antibody

Antibody Specification	
Catalog No.	ITA9202
Product Description	Immunotag™ RPC7 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	RPC7
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:1000-3000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human RPC7
Specificity	RPC7 Antibody detects endogenous levels of total RPC7
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	POLR3G
Accession No.	O15318
Alternate Names	DNA directed RNA polymerase III 32 kDa polypeptide; DNA-directed RNA polymerase III subunit G; DNA-directed RNA polymerase III subunit RPC7; Polr3g; Polymerase (RNA) III (DNA directed) polypeptide G (32kD); RNA polymerase III 32 kDa subunit; RNA polymerase III subunit C7; RPC32; RPC7; RPC7_HUMAN;

Antibody Specification

Description	DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Specific peripheric component of RNA polymerase III which synthesizes small RNAs, such as 5S rRNA and tRNAs (PubMed:20154270). May direct with other members of the RPC3/POLR3C-RPC6/POLR3F-RPC7/POLR3G subcomplex RNA Pol III binding to the TFIIIB-DNA complex via the interactions between TFIIIB and POLR3F. May be involved either in the recruitment and stabilization of the subcomplex within RNA polymerase III, or in stimulating catalytic functions of other subunits during initiation. Plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses. Acts as nuclear and cytosolic DNA sensor involved in innate immune response. Can sense non-self dsDNA that serves as template for transcription into dsRNA. The non-self RNA polymerase III transcripts, such as Epstein-Barr virus-encoded RNAs (EBERs), induce type I interferon and NF- Kappa-B through the RIG-I pathway (PubMed:19609254, PubMed:19631370).
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	26 kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.