

Immunotag™ MARCH8 Antibody

Antibody Specification	
Catalog No.	ITA9225
Product Description	Immunotag™ MARCH8 Antibody
Size	100 µg, 200 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	MARCH8
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:1000-3000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	A synthesized peptide derived from human MARCH8
Specificity	MARCH8 Antibody detects endogenous levels of total 42437
Purification	The antiserum was purified by peptide affinity chromatography.
Form	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt
Gene Name	MARCH8
Accession No.	Q5T0T0
Alternate Names	c MIR; c-MIR; Cellular modulator of immune recognition; E3 ubiquitin protein ligase MARCH8; E3 ubiquitin-protein ligase MARCH8; MARCH 8; MARCH VIII; MARCH-VIII; MARCH8; MARH8_HUMAN; Membrane associated RING CH protein VIII; Membrane associated ring finger (C3HC4) 8; Membrane associated RING finger protein 8; Membrane-associated RING finger protein 8; Membrane-associated RING-CH protein VIII; MIR; OTTHUMP00000058921; RING finger protein 178; RNF178;

Antibody Specification

Description	E3 ubiquitin-protein ligase that mediates ubiquitination of CD86 and MHC class II proteins, such as HLA-DR alpha and beta, and promotes their subsequent endocytosis and sorting to lysosomes via multivesicular bodies. May also promote ubiquitination and endocytosis of TFRC and FAS.
Cell Pathway/ Category	Primary Polyclonal Antibody
Protein MW	33 kDa
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.