

Immunotag™ MEK-4 Monoclonal Antibody

Antibody Specification	
Catalog No.	ITM0437
Product Description	Immunotag™ MEK-4 Monoclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	MEK-4
Clonality	Monoclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,FCM,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/200 - 1/1000. Flow cytometry: 1/200 - 1/400. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Mouse
Immunogen	Purified recombinant fragment of human MEK-4 expressed in E. Coli.
Specificity	MEK-4 Monoclonal Antibody detects endogenous levels of MEK-4 protein.
Purification	Affinity purification
Form	Ascitic fluid containing 0.03% sodium azide.
Gene Name	MAP2K4
Accession No.	P45985 P47809
Alternate Names	MAP2K4; JNKK1; MEK4; MKK4; PRKMK4; SEK1; SERK1; SKK1; Dual specificity mitogen-activated protein kinase kinase 4; MAP kinase kinase 4; MAPKK 4; JNK-activating kinase 1; MAPK/ERK kinase 4; MEK 4; SAPK/ERK kinase 1; SEK1; Stress-activated pro

Antibody Specification

Description	mitogen-activated protein kinase kinase 4(MAP2K4) Homo sapiens This gene encodes a member of the mitogen-activated protein kinase (MAPK) family. Members of this family act as an integration point for multiple biochemical signals and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation, and development. They form a three-tiered signaling module composed of MAPKKKs, MAPKKs, and MAPKs. This protein is phosphorylated at serine and threonine residues by MAPKKKs and subsequently phosphorylates downstream MAPK targets at threonine and tyrosine residues. A similar protein in mouse has been reported to play a role in liver organogenesis. A pseudogene of this gene is located on the long arm of chromosome X. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013],
Cell Pathway/ Category	Regulates Angiogenesis, Stem cell pathway, Regulation of Actin Dynamics, Toll_Like, Cell Growth, ErbB/HER, B Cell Receptor, MAPK_ERK_Growth,MAPK_G_Protein
Protein Expression	Brain,Testis,
Subcellular Localization	nucleus,cytoplasm,cytosol,axon,dendrite cytoplasm,perikaryon,
Protein Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Dual specificity kinase that activates the JUN kinases MAPK8 (JNK1) and MAPK9 (JNK2) as well as MAPK14 (p38) but not MAPK1 (ERK2) or MAPK3 (ERK1).,PTM:Activated by phosphorylation on Ser/Thr by MAP kinase kinase kinases.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase subfamily.,similarity:Contains 1 protein kinase domain.,subunit:Interacts with SPAG9.,tissue specificity:Abundant expression is seen in the skeletal muscle. It is also widely expressed in other tissues.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.