

Immunotag™ PBEF Monoclonal Antibody

| Antibody Specification | |
|------------------------|--|
| Catalog No. | ITM0510 |
| Product Description | Immunotag™ PBEF Monoclonal Antibody |
| Size | 50 µg, 100 µg |
| Conjugation | HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647 |
| IMPORTANT NOTE | This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return. |
| Target Protein | PBEF |
| Clonality | Monoclonal |
| Storage/Stability | -20°C/1 year |
| Application | WB,ELISA |
| Recommended Dilution | Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. |
| Concentration | 1 mg/ml |
| Reactive Species | Human |
| Host Species | Mouse |
| Immunogen | Purified recombinant fragment of PBEF expressed in E. Coli. |
| Specificity | PBEF Monoclonal Antibody detects endogenous levels of PBEF protein. |
| Purification | Affinity purification |
| Form | Ascitic fluid containing 0.03% sodium azide. |
| Gene Name | NAMPT |
| Accession No. | P43490 Q99KQ4 |
| Alternate Names | NAMPT; PBEF; PBEF1; Nicotinamide phosphoribosyltransferase; NAMPRTase; Nampt; Pre-B-cell colony-enhancing factor 1; Pre-B cell-enhancing factor; Visfatin |

Antibody Specification

| | |
|-----------------------------|---|
| Description | nicotinamide phosphoribosyltransferase(NAMPT) Homo sapiens This gene encodes a protein that catalyzes the condensation of nicotinamide with 5-phosphoribosyl-1-pyrophosphate to yield nicotinamide mononucleotide, one step in the biosynthesis of nicotinamide adenine dinucleotide. The protein belongs to the nicotinic acid phosphoribosyltransferase (NAPRTase) family and is thought to be involved in many important biological processes, including metabolism, stress response and aging. This gene has a pseudogene on chromosome 10. [provided by RefSeq, Feb 2011], |
| Cell Pathway/ Category | Nicotinate and nicotinamide metabolism, |
| Protein Expression | Blood,Lung,Skin, |
| Subcellular Localization | extracellular space,nucleoplasm,cytoplasm,cytosol,cell junction,extracellular exosome, |
| Protein Function | catalytic activity:Nicotinamide D-ribonucleotide + diphosphate = nicotinamide + 5-phospho-alpha-D-ribose 1-diphosphate.,caution:Was originally (PubMed:8289818) thought to be a cytokine which acts on early B-lineage precursor cells, by enhancing the effect of IL-7 and SCF on pre-B-cell colony formation.,function:Catalyzes the condensation of nicotinamide with 5-phosphoribosyl-1-pyrophosphate to yield nicotinamide mononucleotide, an intermediate in the biosynthesis of NAD. It is the rate limiting component in the mammalian NAD biosynthesis pathway.,pathway:Cofactor biosynthesis; NAD(+) biosynthesis; nicotinamide ribonucleotide from 5-phospho-alpha-D-ribose 1-diphosphate and nicotinamide: step 1/1.,similarity:Belongs to the NAPRTase family.,tissue specificity:Expressed in large amounts in bone marrow, liver tissue, and muscle. Also present in heart, placenta, lung, and kidney tissues., |
| Usage | For Research Use Only! Not for diagnostic or therapeutic procedures. |