

Immunotag™ TudorSN Monoclonal Antibody

Antibody Specification	
Catalog No.	ITM0635
Product Description	Immunotag™ TudorSN Monoclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	TudorSN
Clonality	Monoclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Mouse
Immunogen	Purified recombinant fragment of TudorSN (aa361-485) expressed in E. Coli.
Specificity	TudorSN Monoclonal Antibody detects endogenous levels of TudorSN protein.
Purification	Affinity purification
Form	Ascitic fluid containing 0.03% sodium azide.
Gene Name	SND1
Accession No.	Q7KZF4 Q78PY7
Alternate Names	SND1; TDRD11; Staphylococcal nuclease domain-containing protein 1; 100 kDa coactivator; EBNA2 coactivator p100; Tudor domain-containing protein 11; p100 co-activator

Antibody Specification

Description	staphylococcal nuclease and tudor domain containing 1(SND1) Homo sapiens This gene encodes a transcriptional co-activator that interacts with the acidic domain of Epstein-Barr virus nuclear antigen 2 (EBNA 2), a transcriptional activator that is required for B-lymphocyte transformation. Other transcription factors that interact with this protein are signal transducers and activators of transcription, STATs. This protein is also thought to be essential for normal cell growth. A similar protein in mammals and other organisms is a component of the RNA-induced silencing complex (RISC). [provided by RefSeq, Jul 2016],
Protein Expression	Brain,Colon carcinoma,Epithelium,Lymph,
Subcellular Localization	nucleus,cytoplasm,mitochondrion,cell-cell adherens junction,membrane,RISC complex,melanosome,extracellular exosome,dense body,
Protein Function	function:Functions as a bridging factor between STAT6 and the basal transcription factor. Plays a role in PIM1 regulation of MYB activity. Functions as a transcriptional coactivator for the Epstein-Barr virus nuclear antigen 2 (EBNA2).,PTM:Phosphorylated by PIM1 in vitro.,sequence caution:The frameshift leads to wrong initiation.,similarity:Contains 1 Tudor domain.,similarity:Contains 4 TNase-like domains.,subcellular location:In IL-4 stimulated cells colocalizes with STAT6 in the nucleus. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.,subunit:Binds to acidic transactivation domain of EBNA2. Interacts with EAV NSP1. Interacts with GTF2E1 and GTF2E2. Forms a ternary complex with STAT6 and POLR2A. Interacts with STAT5.,tissue specificity:Ubiquitously expressed.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.