

Immunotag™ Vimentin Monoclonal Antibody

Antibody Specification	
Catalog No.	ITM0645
Product Description	Immunotag™ Vimentin Monoclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Vimentin
Clonality	Monoclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Mouse
Immunogen	Purified recombinant fragment of Vimentin expressed in E. Coli.
Specificity	Vimentin Monoclonal Antibody detects endogenous levels of Vimentin protein.
Purification	Affinity purification
Form	Ascitic fluid containing 0.03% sodium azide.
Gene Name	VIM
Accession No.	P08670 P20152
Alternate Names	VIM; Vimentin

Antibody Specification

Description	vimentin(VIM) Homo sapiens This gene encodes a member of the intermediate filament family. Intermediate filamentents, along with microtubules and actin microfilaments, make up the cytoskeleton. The protein encoded by this gene is responsible for maintaining cell shape, integrity of the cytoplasm, and stabilizing cytoskeletal interactions. It is also involved in the immune response, and controls the transport of low-density lipoprotein (LDL)-derived cholesterol from a lysosome to the site of esterification. It functions as an organizer of a number of critical proteins involved in attachment, migration, and cell signaling. Mutations in this gene causes a dominant, pulverulent cataract.[provided by RefSeq, Jun 2009],
Protein Expression	Adipose tissue,Brain,Cajal-Retzius cell,Cervix,Coronary artery,Embryo,Epithelium,Fetal brain cortex
Subcellular Localization	cytoplasm,peroxisome,cytosol,cytoskeleton,intermediate filament,plasma membrane,focal adhesion,extracellular matrix,cell leading edge,neuron projection,intermediate filament cytoskeleton,extracellular exosome,
Protein Function	function:Vimentins are class-III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells.,online information:Vimentin entry,PTM:One of the most prominent phosphoproteins in various cells of mesenchymal origin. Phosphorylation is enhanced during cell division, at which time vimentin filaments are significantly reorganized.,sequence caution:Intron retention.,similarity:Belongs to the intermediate filament family.,subunit:Homopolymer. Interacts with HCV core protein. Interacts with LGSN and SYNM.,tissue specificity:Highly expressed in fibroblasts, some expression in T- and B-lymphocytes, and little or no expression in Burkitt's lymphoma cell lines. Expressed in many hormone-independent mammary carcinoma cell lines.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.