

Immunotag™ PSR Monoclonal Antibody

Antibody Specification	
Catalog No.	ITM1087
Product Description	Immunotag™ PSR Monoclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PSR
Clonality	Monoclonal
Storage/Stability	-20°C/1 year
Application	WB,IF
Recommended Dilution	Western Blot: 1/1000 - 1/2000. Immunofluorescence: 1/100 - 1/500. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat,Bovine,Dog
Host Species	Mouse
Immunogen	Purified recombinant human PSR (N-terminus) protein fragments expressed in Ecoli
Specificity	PSR Monoclonal Antibody detects endogenous levels of PSR protein.
Purification	Affinity purification
Form	Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.2% sodium azide, 50% glycerol.
Gene Name	JMJD6
Accession No.	Q6NYC1 Q9ERI5 Q6AYK2
Alternate Names	JMJD6; KIAA0585; PTDSR; Bifunctional arginine demethylase and lysyl-hydroxylase JMJD6; Histone arginine demethylase JMJD6; JmjC domain-containing protein 6; Jumonji domain-containing protein 6; Lysyl-hydroxylase JMJD6; Peptide-lysine 5-diox

Antibody Specification

Description	arginine demethylase and lysine hydroxylase(JMJD6) Homo sapiens This gene encodes a nuclear protein with a JmjC domain. JmjC domain-containing proteins are predicted to function as protein hydroxylases or histone demethylases. This protein was first identified as a putative phosphatidylserine receptor involved in phagocytosis of apoptotic cells; however, subsequent studies have indicated that it does not directly function in the clearance of apoptotic cells, and questioned whether it is a true phosphatidylserine receptor. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],
Protein Expression	Brain,Skin,Stomach cancer,Uterus,
Subcellular Localization	nucleus,nucleoplasm,nucleolus,cytosol,plasma membrane,intracellular ribonucleoprotein complex,
Protein Function	caution:Was initially thought to constitute the phosphatidylserine receptor, a receptor that mediates recognition of phosphatidylserine, a specific marker only present at the surface of apoptotic cells. Phosphatidylserine receptor probably participates in apoptotic cell phagocytosis. This protein was identified using phage display expressing mAb 217, an antibody that specifically recognizes phosphatidylserine receptor. However, its nuclear localization and the fact that mAb 217 antibody still recognizes the phosphatidylserine receptor in mice lacking JMJD6, strongly suggest that it does not constitute the receptor for phosphatidylserine and is not involved in apoptotic cell removal.,domain:The nuclear localization signal motifs are necessary and sufficient to target it into the nucleus.,function:Arginine demethylase which demethylates histone H3 at 'Arg-2' (H3R2me) and histone H4 at 'Arg-3' (H4R3me). Required for differentiation of multiple organs during embryogenesis. Probably acts as a key regulator of hematopoietic differentiation. Seems to be necessary for the regulation of macrophage cytokine responses.,induction:Up-regulated upon cytokine treatment, but not upon TNF-alpha treatment.,similarity:Belongs to the PTDSR family.,similarity:Contains 1 JmjC domain.,tissue specificity:Highly expressed in the heart, skeletal muscle and kidney. Expressed at moderate or low level in brain, placenta, lung, liver, pancreas, spleen, thymus, prostate, testis and ovary. Up-regulated in many patients with chronic pancreatitis. Expressed in nursing thymic epithelial cells.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.