

Immunotag™ SMAD5(C-term) mouse mAb

Antibody Specification	
Catalog No.	ITM1259
Product Description	Immunotag™ SMAD5(C-term) mouse mAb
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	SMAD5 (C-term)
Clonality	Monoclonal
Storage/Stability	-20°C/1 year
Application	WB,ICC,FC
Recommended Dilution	wb 1:1000 icc 1:75 fcm 1:100
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Mouse
Immunogen	Purified recombinant human SMAD5 (C-terminus) protein fragments expressed in E.coli.
Specificity	This antibody detects endogenous levels of SMAD5 (C-terminus) and does not cross-react with related proteins.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	smad5
Accession No.	Q99717 P97454
Alternate Names	DKFZp781C1895;DKFZp781O1323;Dwfc;hSmad 5;hSmad5;JV5 1;JV5-1;MAD homolog 5;MAD mothers against decapentaplegic homolog 5;MAD, mothers against decapentaplegic homolog 5;MADH 5;MADH5;Mothers against decapentaplegic homolog 5;Mothers against DPP homolog 5;MusMLP; SMA and MAD related protein 5;SMAD 5;SMAD family member 5;SMAD mothers against DPP homolog 5;Smad5;Smad5;SMAD5_HUMAN.

Antibody Specification

Description	SMAD family member 5(SMAD5) Homo sapiens The protein encoded by this gene is involved in the transforming growth factor beta signaling pathway that results in an inhibition of the proliferation of hematopoietic progenitor cells. The encoded protein is activated by bone morphogenetic proteins type 1 receptor kinase, and may be involved in cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2014],
Cell Pathway/ Category	TGF-beta,
Protein Expression	Colon carcinoma,Epithelium,Testis,Trachea,Uterus,
Subcellular Localization	intracellular,nucleus,nucleoplasm,transcription factor complex,cytoplasm,cytosol,integral component of membrane,protein complex,SMAD protein complex,
Protein Function	function:Transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase. SMAD5 is a receptor-regulated SMAD (R-SMAD).,PTM:Phosphorylated on serine by BMP (bone morphogenetic proteins) type 1 receptor kinase.,PTM:Ubiquitin-mediated proteolysis by SMAD-specific E3 ubiquitin ligase SMURF1.,similarity:Belongs to the dwarfin/SMAD family.,similarity:Contains 1 MH1 (MAD homology 1) domain.,similarity:Contains 1 MH2 (MAD homology 2) domain.,subcellular location:Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4.,subunit:May form trimers with the co-SMAD SMAD4. Interacts with PEBP2-alpha subunit and SMURF1. Interacts with SUV39H1 and SUV39H2.,tissue specificity:Ubiquitous.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.