

Immunotag™ Transferrin mouse mAb

Antibody Specification	
Catalog No.	ITM1463
Product Description	Immunotag™ Transferrin mouse mAb
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Transferrin
Clonality	Monoclonal
Storage/Stability	-20°C/1 year
Application	ELISA
Recommended Dilution	ELISA 1:10000-20000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Mouse
Immunogen	Transferrin human.
Specificity	This antibody detects Human Transferrin.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	transferrin
Accession No.	P02787 Q921I1
Alternate Names	Apotransferrin; Beta 1 metal binding globulin; Beta-1 metal-binding globulin; DKFZp781D0156; PRO1400; PRO1557; PRO2086; Serotransferrin; Serotransferrin precursor; Siderophilin; TF; TFQTL1; Transferin; Transferrin; TRFE_HUMAN.

Antibody Specification

Description	transferrin(TF) Homo sapiens This gene encodes a glycoprotein with an approximate molecular weight of 76.5 kDa. It is thought to have been created as a result of an ancient gene duplication event that led to generation of homologous C and N-terminal domains each of which binds one ion of ferric iron. The function of this protein is to transport iron from the intestine, reticuloendothelial system, and liver parenchymal cells to all proliferating cells in the body. This protein may also have a physiologic role as granulocyte/pollen-binding protein (GPBP) involved in the removal of certain organic matter and allergens from serum. [provided by RefSeq, Sep 2009],
Protein Expression	Bile,Brain,Cajal-Retzius cell,Caudate nucleus,Fetal brain cortex,Fetal liver,Heart,Liver,Plasma,Sal
Subcellular Localization	extracellular region,extracellular space,early endosome,late endosome,clathrin-coated pit,basal plasma membrane,cell surface,endosome membrane,cytoplasmic, membrane-bounded vesicle,apical plasma membrane,vesicle coat,endocytic vesicle,
Protein Function	disease:Defects in TF are the cause of atransferrinemia [MIM:209300]. Atransferrinemia is rare autosomal recessive disorder characterized by iron overload and hypochromic anemia.,function:Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with the binding of an anion, usually bicarbonate. It is responsible for the transport of iron from sites of absorption and heme degradation to those of storage and utilization. Serum transferrin may also have a further role in stimulating cell proliferation.,online information:Transferrin entry,polymorphism:Different polymorphic variants of transferrin are known. The sequence shown is the predominant electrophoretic variant (C1 or TF*C1).,similarity:Belongs to the transferrin family.,similarity:Contains 2 transferrin-like domains.,subunit:Monomer.,tissue specificity:Expressed by the liver and secreted in plasma.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.