

Immunotag™ VE-Cadherin mouse Monoclonal Antibody(8E5)

Antibody Specification	
Catalog No.	ITM3673
Product Description	Immunotag™ VE-Cadherin mouse Monoclonal Antibody(8E5)
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	VE-Cadherin (8E5)
Clonality	Monoclonal
Storage/Stability	-20°C/1 year
Application	IHC-p
Recommended Dilution	IHC 1:100-200
Concentration	1 mg/ml
Reactive Species	Human,Rat,Mouse
Host Species	Mouse
Immunogen	Synthetic Peptide of VE-Cadherin at AA range of 670-750
Specificity	VE-Cadherin protein detects endogenous levels of CDH5
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	CDH5
Accession No.	P33151 P55284
Alternate Names	CDH5

Antibody Specification

Description	cadherin 5(CDH5) Homo sapiens This gene encodes a classical cadherin of the cadherin superfamily. The encoded preproprotein is proteolytically processed to generate the mature glycoprotein. This calcium-dependent cell-cell adhesion molecule is comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Functioning as a classical cadherin by imparting to cells the ability to adhere in a homophilic manner, this protein plays a role in endothelial adherens junction assembly and maintenance. This gene is located in a gene cluster in a region on the long arm of chromosome 16 that is involved in loss of heterozygosity events in breast and prostate cancer. [provided by RefSeq, Nov 2015],
Cell Pathway/ Category	Cell adhesion molecules (CAMs),Leukocyte transendothelial migration,
Protein Expression	Aorta endothelial cell,Brain,Endothelial cell,PCR rescued clones,Placenta,Plasma,
Subcellular Localization	plasma membrane,cell-cell junction,adherens junction,bicellular tight junction,external side of plasma membrane,cell surface,membrane,integral component of membrane,cell junction,
Protein Function	function:Cadherins are calcium dependent cell adhesion proteins.,function:Cadherins are calcium dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. This cadherin may play a important role in endothelial cell biology through control of the cohesion and organization of the intercellular junctions. It associates with alpha-catenin forming a link to the cytoskeleton.,similarity:Contains 5 cadherin domains.,subcellular location:Found at cell-cell boundaries and probably at cell-matrix boundaries.,tissue specificity:Endothelial tissues and brain.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.