

Immunotag™ PXDC2 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN0125
Product Description	Immunotag™ PXDC2 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PXDC2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein . at AA range: 50-130
Specificity	PXDC2 Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	PLXDC2 TEM7R UNQ2514/PRO6003
Accession No.	Q6UX71 Q9DC11
Description	function:May play a role in tumor angiogenesis.,similarity:Belongs to the plexin family.,similarity:Contains 1 PSI domain.,subunit:Interacts with CTTN.,tissue specificity:Expressed in the endothelial cells of the stroma but not in the endothelial cells of normal colonic tissue.,
Protein Expression	Liver,Ovary,Plasma,

Antibody Specification

Subcellular Localization	integral component of membrane,extracellular exosome,
Protein Function	function:May play a role in tumor angiogenesis.,similarity:Belongs to the plexin family.,similarity:Contains 1 PSI domain.,subunit:Interacts with CTTN.,tissue specificity:Expressed in the endothelial cells of the stroma but not in the endothelial cells of normal colonic tissue.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.