

Immunotag™ RBPJL Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN0154
Product Description	Immunotag™ RBPJL Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	RBPJL
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein . at AA range: 140-220
Specificity	RBPJL Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	RBPJL RBPL RBPSUHL
Accession No.	Q9UBG7 O08674

Antibody Specification

Description	recombination signal binding protein for immunoglobulin kappa J region like(RBPJL) Homo sapiens This gene encodes a member of the suppressor of hairless protein family. A similar protein in mouse is a transcription factor that binds to DNA sequences almost identical to that bound by the Notch receptor signaling pathway transcription factor recombining binding protein J. The mouse protein has been shown to activate transcription in concert with Epstein-Barr virus nuclear antigen-2. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013],
Cell Pathway/ Category	Notch,
Subcellular Localization	nucleus,transcription factor complex,
Protein Function	function:Putative transcription factor, which cooperates with EBNA2 to activate transcription.,similarity:Belongs to the Su(H) family.,similarity:Contains 1 IPT/TIG domain.,subunit:Interacts weakly with EBNA2. Does not interact with any Notch proteins.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.