

Immunotag™ RFOX1 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN0464
Product Description	Immunotag™ RFOX1 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	RFOX1
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein . at AA range: 50-130
Specificity	RFOX1 Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	RBFOX1 A2BP A2BP1 FOX1 HRNBP1
Accession No.	Q9NWB1 Q9JJ43

## Antibody Specification

Description	RNA binding protein, fox-1 homolog 1(RBFOX1) Homo sapiens The Fox-1 family of RNA-binding proteins is evolutionarily conserved, and regulates tissue-specific alternative splicing in metazoa. Fox-1 recognizes a (U)GCAUG stretch in regulated exons or in flanking introns. The protein binds to the C-terminus of ataxin-2 and may contribute to the restricted pathology of spinocerebellar ataxia type 2 (SCA2). Ataxin-2 is the product of the SCA2 gene which causes familial neurodegenerative diseases. Fox-1 and ataxin-2 are both localized in the trans-Golgi network. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2011],
Protein Expression	Brain,
Subcellular Localization	nucleus,cytoplasm,trans-Golgi network,
Protein Function	function:RNA-binding protein that regulates alternative splicing events by binding to 5'-UGCAUGU-3' elements. Regulates alternative splicing of tissue-specific exons and of differentially spliced exons during erythropoiesis.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Binds to the C-terminal of ATXN2.,tissue specificity:Predominantly expressed in muscle and brain.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.