

Immunotag™ PDE1A Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN0520
Product Description	Immunotag™ PDE1A Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PDE1A
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	PDE1A Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	PDE1A
Accession No.	P54750 Q61481

Antibody Specification

Description	phosphodiesterase 1A(PDE1A) Homo sapiens Cyclic nucleotide phosphodiesterases (PDEs) play a role in signal transduction by regulating intracellular cyclic nucleotide concentrations through hydrolysis of cAMP and/or cGMP to their respective nucleoside 5-prime monophosphates. Members of the PDE1 family, such as PDE1A, are Ca(2+)/calmodulin (see CALM1; MIM 114180)-dependent PDEs (CaM-PDEs) that are activated by calmodulin in the presence of Ca(2+) (Michibata et al., 2001 [PubMed 11342109]; Fidock et al., 2002 [PubMed 11747989]).[supplied by OMIM, Oct 2009],
Cell Pathway/ Category	Purine metabolism,Calcium,Taste transduction,
Protein Expression	Brain,Cerebral cortex,Heart,Lung,Uterus,
Subcellular Localization	nucleus,cytosol,neuronal cell body,
Protein Function	catalytic activity:Nucleoside 3',5'-cyclic phosphate + H(2)O = nucleoside 5'-phosphate.,enzyme regulation:Type I PDE are activated by the binding of calmodulin in the presence of Ca(2+).,function:Has a higher affinity for cGMP than for cAMP.,similarity:Belongs to the cyclic nucleotide phosphodiesterase family.,subunit:Homodimer.,tissue specificity:Several tissues, including brain, kidney, testes and heart.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.