Immunotag™ RPOM Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN0665
Product Description	Immunotag™ RPOM Polyclonal Antibody
Size	50 μg, 100 μg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	RPOM
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	RPOM Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	POLRMT
Accession No.	O00411 Q8BKF1

Antibody Specification	
Description	RNA polymerase mitochondrial(POLRMT) Homo sapiens This gene encodes a mitochondrial DNA-directed RNA polymerase. The gene product is responsible for mitochondrial gene expression as well as for providing RNA primers for initiation of replication of the mitochondrial genome. Although this polypeptide has the same function as the three nuclear DNA-directed RNA polymerases, it is more closely related to RNA polymerases of phage and mitochondrial polymerases of lower eukaryotes. [provided by RefSeq, Jul 2008],
Protein Expression	Aorta endothelial cell,Brain,Liver,
Subcellular Localization	mitochondrion,mitochondrial matrix,mitochondrial nucleoid,
Protein Function	catalytic activity: Nucleoside triphosphate $+$ RNA(n) = diphosphate $+$ RNA(n+1)., function: DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates., similarity: Belongs to the phage and mitochondrial RNA polymerase family., subunit: Interacts with TFB1M and TFB2M, leading to the stimulation of transcription.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.

www.gbiosciences.com

© 2018 Geno Technology Inc., USA. All Rights Reserved.