

Immunotag™ MYO5C Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN0908
Product Description	Immunotag™ MYO5C Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	MYO5C
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein . at AA range: 850-930
Specificity	MYO5C Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	MYO5C
Accession No.	Q9NQX4
Description	function:May be involved in transferrin trafficking. Likely to power actin-based membrane trafficking in many physiologically crucial tissues.,similarity:Contains 1 dilute domain.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 6 IQ domains.,tissue specificity:Expressed chiefly in non-neuronal tissues. Particularly abundant in epithelial and glandular tissues including pancreas, prostate, mammary, stomach, colon and lung.,
Protein Expression	Cord blood,Hepatoma,Peripheral Nervous System,

Antibody Specification

Subcellular Localization	myosin complex,extracellular exosome,
Protein Function	function:May be involved in transferrin trafficking. Likely to power actin-based membrane trafficking in many physiologically crucial tissues.,similarity:Contains 1 dilute domain.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 6 IQ domains.,tissue specificity:Expressed chiefly in non-neuronal tissues. Particularly abundant in epithelial and glandular tissues including pancreas, prostate, mammary, stomach, colon and lung.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.