

Immunotag™ NRX1B Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN0951
Product Description	Immunotag™ NRX1B Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	NRX1B
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Rat,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein, at AA range: 260-340
Specificity	NRX1B Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	NRXN1
Accession No.	P58400 P0DI97 Q63373

Antibody Specification

Description	<p>alternative products:A number of isoforms are produced by alternative promoter usage including the alpha-type (shown here) and beta-type (AC P58400) isoforms which differ in their N-terminus. Additional isoforms may be produced by alternative splicing,alternative products:A number of isoforms, alpha-type (AC Q9ULB1) and beta-type (shown here), are produced by alternative promoter usage. Beta-type isoforms differ from alpha-type isoforms in their N-terminus,function:Neuronal cell surface protein that may be involved in cell recognition and cell adhesion by forming intracellular junctions through binding to neuroligins. May play a role in formation or maintenance of synaptic junctions. May mediate intracellular signaling.,function:Neuronal cell surface protein that may be involved in cell recognition and cell adhesion. May mediate intracellular signaling.,PTM:Highly O-glycosylated and minor N-glycosylated.,PTM:N- and O-glycosylated.,similarity:Belongs to the neurexin family.,similarity:Contains 1 laminin G-like domain.,similarity:Contains 3 EGF-like domains.,similarity:Contains 6 laminin G-like domains.,subunit:The cytoplasmic C-terminal region binds to CASK, CASKIN1 and APBA1. The laminin G-like domain 2 binds to NXPH1. Specific isoforms bind to alpha-dystroglycan and to alpha-latroxin. Interacts with SYT13 and SYTL1.,subunit:The cytoplasmic C-terminal region binds to CASK. Isoforms Beta 4b bind neuroligins NLGN1, NLGN2 and NLGN3, alpha-dystroglycan and alpha-latrotoxin.,tissue specificity:Heart and brain.,</p>
Cell Pathway/ Category	Cell adhesion molecules (CAMs),
Protein Function	<p>A number of isoforms are produced by alternative promoter usage including the alpha-type (shown here) and beta-type (AC P58400) isoforms which differ in their N-terminus. Additional isoforms may be produced by alternative splicing,A number of isoforms, alpha-type (AC Q9ULB1) and beta-type (shown here), are produced by alternative promoter usage. Beta-type isoforms differ from alpha-type isoforms in their N-terminus,function:Neuronal cell surface protein that may be involved in cell recognition and cell adhesion by forming intracellular junctions through binding to neuroligins. May play a role in formation or maintenance of synaptic junctions. May mediate intracellular signaling.,function:Neuronal cell surface protein that may be involved in cell recognition and cell adhesion. May mediate intracellular signaling.,PTM:Highly O-glycosylated and minor N-glycosylated.,PTM:N- and O-glycosylated.,similarity:Belongs to the neurexin family.,similarity:Contains 1 laminin G-like domain.,similarity:Contains 3 EGF-like domains.,similarity:Contains 6 laminin G-like domains.,subunit:The cytoplasmic C-terminal region binds to CASK, CASKIN1 and APBA1. The laminin G-like domain 2 binds to NXPH1. Specific isoforms bind to alpha-dystroglycan and to alpha-latroxin. Interacts with SYT13 and SYTL1.,subunit:The cytoplasmic C-terminal region binds to CASK. Isoforms Beta 4b bind neuroligins NLGN1, NLGN2 and NLGN3, alpha-dystroglycan and alpha-latrotoxin.,tissue specificity:Heart and brain.,</p>
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.