

Immunotag™ PEX12 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1039
Product Description	Immunotag™ PEX12 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PEX12
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Rat,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein, at AA range: 180-260
Specificity	PEX12 Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	PEX12 PAF3
Accession No.	O00623 Q8VC48 O88177

Antibody Specification

Description	<p>peroxisomal biogenesis factor 12(PEX12) Homo sapiens This gene belongs to the peroxin-12 family. Peroxins (PEXs) are proteins that are essential for the assembly of functional peroxisomes. The peroxisome biogenesis disorders (PBDs) are a group of genetically heterogeneous autosomal recessive, lethal diseases characterized by multiple defects in peroxisome function. The peroxisomal biogenesis disorders are a heterogeneous group with at least 14 complementation groups and with more than 1 phenotype being observed in cases falling into particular complementation groups. Although the clinical features of PBD patients vary, cells from all PBD patients exhibit a defect in the import of one or more classes of peroxisomal matrix proteins into the organelle. Defects in this gene are a cause of Zellweger syndrome (ZWS). [provided by RefSeq, Oct 2008],</p>
Protein Expression	Fetal brain,Testis,
Subcellular Localization	peroxisome,peroxisomal membrane,integral component of peroxisomal membrane,peroxisomal importomer complex,
Protein Function	<p>disease:Defects in PEX12 are a cause of Zellweger syndrome (ZWS) [MIM:214100]. ZWS is a fatal peroxisome biogenesis disorder characterized by dysmorphic facial features, hepatomegaly, ocular abnormalities, renal cysts, hearing impairment, profound psychomotor retardation, severe hypotonia and neonatal seizures. Death occurs within the first year of life.,disease:Defects in PEX12 are the cause of peroxisome biogenesis disorder complementation group 3 (PBD-CG3) [MIM:601758]. PBD refers to a group of peroxisomal disorders arising from a failure of protein import into the peroxisomal membrane or matrix. The PBD group is comprised of four disorders: Zellweger syndrome (ZWS), neonatal adrenoleukodystrophy (NALD), infantile Refsum disease (IRD), and classical rhizomelic chondrodysplasia punctata (RCDP). ZWS, NALD and IRD are distinct from RCDP and constitute a clinical continuum of overlapping phenotypes known as the Zellweger spectrum. The PBD group is genetically heterogeneous with at least 14 distinct genetic groups as concluded from complementation studies.,function:Required for protein import into peroxisomes.,similarity:Belongs to the pex2/pex10/pex12 family.,similarity:Contains 1 RING-type zinc finger.,subunit:Interacts with PEX5 and PEX10. Interacts with PEX19 via its cytoplasmic domain.,</p>
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.