

Immunotag™ PIGQ Polyclonal Antibody

Antibody Specification	
Catalog No.	ITN1046
Product Description	Immunotag™ PIGQ Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PIGQ
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human protein . at AA range: 130-210
Specificity	PIGQ Polyclonal Antibody detects endogenous levels of protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Gene Name	PIGQ GPI1
Accession No.	Q9BRB3 Q9QYT7

Antibody Specification

Description	phosphatidylinositol glycan anchor biosynthesis class Q(PIGQ) Homo sapiens This gene is involved in the first step in glycosylphosphatidylinositol (GPI)-anchor biosynthesis. The GPI-anchor is a glycolipid found on many blood cells and serves to anchor proteins to the cell surface. This gene encodes a N-acetylglucosaminyl transferase component that is part of the complex that catalyzes transfer of N-acetylglucosamine (GlcNAc) from UDP-GlcNAc to phosphatidylinositol (PI). Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jun 2012],
Cell Pathway/ Category	Glycosylphosphatidylinositol(GPI)-anchor biosynthesis,
Protein Expression	Melanoma,Retinoblastoma,
Subcellular Localization	glycosylphosphatidylinositol-N-acetylglucosaminyltransferase (GPI-GnT) complex,endoplasmic reticulum membrane,integral component of membrane,
Protein Function	catalytic activity:UDP-N-acetyl-D-glucosamine + 1-phosphatidyl-1D-myo-inositol = UDP + 6-(N-acetyl-alpha-D-glucosaminyl)-1-phosphatidyl-1D-myo-inositol.,function:Part of the complex catalyzing the transfer of N-acetylglucosamine from UDP-N-acetylglucosamine to phosphatidylinositol, the first step of GPI biosynthesis.,pathway:Glycolipid biosynthesis; glycosylphosphatidylinositol-anchor biosynthesis.,similarity:Belongs to the PIGQ family.,subunit:Associates with PIGA, PIGC, PIGH, PIGP and DPM2. The latter is not essential for activity.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.